

STOCKPORT SAFEGUARDING ADULTS BOARD

01 Background

Protection from fire and prevention of future deaths: Several tragic deaths of residents within care homes have led to the Coroner making recommendations about the appropriate levels of fire safety within care facilities.

Are these deaths avoidable? – Yes. The issues need urgent consideration and action by care home Registered Managers, nominated individuals and care teams, to identify critical risks associated with individuals.

The Fire Safety Order 2005 requires the identification of individuals at risk as part of the fire safety risk assessment for the premises and to take appropriate action to remove or reduce the risk.

07 Question

Is the person a smoker?
Are emollients being applied? Does this product contain paraffin? If Yes?
Share the risks with:
Individuals
GP/Nurse Prescriber
Family member
CONSIDER AN ALTERNATIVE PRODUCT

02 Why it matters

A personal risk assessment for each resident is critical for their own safety and that of other residents and staff. This will assess the needs of the individual in conjunction with care workers and family and consider their habits, physical and mental capacity, and their environment. The risk assessment should be recorded and considered as part of their care plan, other assessments and personal evacuation plans, and kept under review.

03 Information

Coroner's advice is that you must consider the risk posed by individuals smoking on your premises, particularly if the person at risk has limited mobility. This follows inquests into the deaths of high-risk smokers with mobility problems from burn injuries as a result of matches or cigarettes dropping on to clothing or bedding.

04 Emollients

Emollient creams are used to treat dry skin conditions and some patients are often in bed for lengthy periods due to illness or impaired mobility. Certain creams can be highly flammable (e.g. those that are paraffin-based) and pose a significant risk in the event of a fire. The fire risk posed by the use of such emollient creams is significantly increased when the user is smoking. The individual's bedding and clothing can also become impregnated with cream, increasing flammability and their risk to the user.

Emollients & Smoking

05 Information

Following recent fire deaths the Coroner highlighted that the use of such creams should be risk assessed and action taken to reduce the harm. The use of emollient creams must be considered in your fire risk assessment to ensure that all reasonably practicable steps are taken to reduce the risk of a fire and its likelihood of occurring.

<http://www.nrls.npsa.nhs.uk/patientsafety-videos/paraffin-based-skinproducts/>

06 What to do

This increased risk of fire posed by smoking whilst using flammable emollient creams is so significant that it must be avoided.

Fire retardant covers, bedding or clothing for smokers must always be provided.

There must be sufficient numbers for items to be laundered at the correct temperature.

This is the responsibility of the care home owing a duty of care for the health, safety and wellbeing of individuals who may be at heightened risk.



STOCKPORT
METROPOLITAN BOROUGH COUNCIL

Safeguarding
Adults
in Stockport